

F3342(c) US

1

FROZEN CONFECTIONERY PRODUCT

Field of the invention

The present invention relates to frozen water ice
5 confectionery products which comprise a plurality of
individual confections and which contain ice structuring
proteins.

Background to the invention

10 The frozen confectionery industry is constantly seeking to
devise novel products that will appeal to consumers. Whilst
frozen confectionery products, such as ice cream and water
ices tend to be sold either in containers, e.g. tubs of ice
cream or cartridge dispensers, or as individually wrapped
15 items such as ice lollies/popsicles, a relatively recent
product innovation is in the form of single serve containers
filled with a plurality of water ice beads. The beads are
manufactured by a process which involves feeding uniformly
sized drops of a liquid composition into a freezing chamber,
20 typically filled with liquid nitrogen.

Water ices typically have from 25 to 35 wt% total solids, of
which a large proportion is sugar. However, it has been
found that if the solids content is above about 6 wt%, then
25 the beads are too soft and tend to stick together and
sinter, as well as deform. Unfortunately this means that it
is difficult to add ingredients such as sugar that would
produce the taste preferred by consumers since the amount of
sugar needed raises the total amount of solids above the
30 desired level. Accordingly, it has been necessary to use
artificial sweeteners, which is less preferred by consumers,
for reasons of taste as well as concerns about the use of
artificial sweeteners.

Summary of the invention

We have now found that the addition of ice structuring proteins to frozen water ice products reduces their tendency to stick and allows the production of free flowing frozen confectionery products that maintain their free-flowing characteristics for longer and at higher storage temperatures than existing products. The appearance of such products is significantly improved compared to existing products even after storage at temperatures above about -20°C for several weeks.

Furthermore, it has been shown to be possible to increase the total solids content significantly above that for existing free-flowing products that lack ISPs without adversely affecting the free-flowing characteristics following storage. This has enabled the inclusion of effective amounts of sugar which in turn improves the taste and flavour of the product.

Accordingly, the present invention provides a frozen confectionery product comprising a plurality of discrete water ice confections, each discrete water ice confection being able to contact directly other discrete water ice confections in the product, which water ice confections comprise an ice structuring protein (ISP), at least 6 wt% solids and have an average volume of less than 1 ml.

Preferably the product comprises at least 10 discrete water ice confections, such as at least 20, 50 or 100 discrete water ice confections.

In a preferred embodiment the discrete water ice confections have an average volume of less than 0.5 ml. The frozen confections may, for example, be in the form of beads.

5 In a related aspect, the present invention provides a product comprising a container filled with a frozen confectionery product of the invention. Preferably the container has a volume of from 100 ml to 1000 ml.

10 The present invention also provides a retail unit comprising a plurality of containers, each container comprising a product of the invention wherein the product in each container is different.

15 Detailed description of the invention

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art (e.g. in frozen confectionery manufacture, molecular biology and
20 biochemistry). Definitions and descriptions of various terms and techniques used in frozen confectionery manufacture are found in Ice Cream, 4th Edition, Arbuckle (1986), Van Nostrand Reinhold Company, New York, NY. Standard techniques are used for molecular and biochemical methods
25 (see generally, Sambrook et al., Molecular Cloning: A Laboratory Manual, 3rd ed. (2001) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. and Ausubel et al., Short Protocols in Molecular Biology (1999) 4th Ed, John Wiley & Sons, Inc. - and the full version entitled
30 Current Protocols in Molecular Biology).

Ice structuring proteins

Ice structuring proteins (ISPs) are proteins that can influence the shape and size of the crystals of ice formed when freezing does occur, and inhibit recrystallisation of ice (Clarke et al., 2002, Cryoletters 23: 89-92). Many of these proteins were identified originally in organisms that live in sub-zero environments and are thought to protect the organism from the deleterious effects of the formation of ice crystals in the cells of the organism. For this reason many ice structuring proteins are also known as antifreeze proteins (AFPs). In the context of the present invention, an ISP is defined as a protein that has ice recrystallisation inhibitory (RI) activity.

Ice recrystallisation inhibitory activity properties can conveniently be measured by means of a modified splat assay as described in WO 00/53029.

2.5 μ l of the solution under investigation in 30% (w/w) sucrose is transferred onto a clean, appropriately labelled, 16 mm circular coverslip. A second coverslip is placed on top of the drop of solution and the sandwich pressed together between finger and thumb. The sandwich is dropped into a bath of hexane held at -80°C in a box of dry ice. When all sandwiches have been prepared, sandwiches are transferred from the -80°C hexane bath to the viewing chamber containing hexane held at -6°C using forceps pre-cooled in the dry ice. Upon transfer to -6°C , sandwiches can be seen to change from a transparent to an opaque appearance. Images are recorded by video camera and grabbed into an image analysis system (LUCIA, Nikon) using a 20x objective. Images of each splat are recorded at time = 0 and again after 60 minutes. The size of the ice-crystals in both

assays is compared by placing the slides within a temperature controlled cryostat cabinet (Bright Instrument Co Ltd, Huntington, UK). Images of the samples are transferred to a Quantimet 520 MC image analysis system
5 (Leica, Cambridge UK) by means of a Sony monochrome CCD videocamera.

Ice crystal sizing can be performed by hand-drawing around the ice-crystals. Typically, at least 100 to 400 crystals
10 are sized for each sample. The ice crystal size is taken as being the longest dimension of the 2D projection of each crystal. The average crystal size is determined as the number average of the individual crystal sizes. The size of the ice-crystals in both assays is compared. If the size at
15 30-60 minutes is similar or only moderately (less than 10%) increased compared to the size at $t=0$, and/or the crystal size is less than 20 micrometer, preferably from 5 to 15 micrometer this is an indication of good ice-crystal recrystallisation properties.

20

Significant ice recrystallisation inhibitory activity can be defined as where a 0.01 wt% solution of the ISP in 30 wt% sucrose, cooled rapidly (at least $\Delta 50^{\circ}\text{C}$ per minute) to -40°C , heated rapidly (at least $\Delta 50^{\circ}\text{C}$ per minute) to -6°C and
25 then held at this temperature results in an increase in average ice crystal size over one hour of less than 5 μm .

Types of ISPs

ISPs for use according to the present invention can be
30 derived from any source provided they are suitable for inclusion in food products. ISPs have been identified to date in fish, plants, lichen, fungi, micro-organisms and

insects. In addition, a number of synthetic ISPs have been described.

Examples of fish ISP materials are AFGP (for example
5 obtainable from Atlantic cod, Greenland cod and Tomcod),
Type I ISP (for example obtainable from Winter flounder,
Yellowtail flounder, Shorthorn sculpin and Grubby sculpin),
Type II ISP (for example obtainable from Sea raven, Smelt
and Atlantic herring) and Type III ISP (for example
10 obtainable from Ocean pout, Atlantic wolffish, Radiated
shanny, Rock gunnel and Laval's eelpout).

Type III ISPs are particularly preferred. Type III ISPs
typically have a molecular weight of from about 6.5 to about
15 14 kDa, a beta sandwich secondary structure and a globular
tertiary structure. A number of genes encoding type III ISPs
have been cloned (Davies and Hew, 1990, FASEB J. 4: 2460-
2468). A particularly preferred type III ISP is type III
HPLC-12 (Accession No. P19614 in the Swiss-Prot protein
20 database).

Lichen AFPs are described in WO99/37673 and WO01/83534.

Examples of plants in which ISPs have been obtained are
25 described in WO 98/04699 and WO 98/4148 and include garlic-
mustard, blue wood aster, spring oat, winter cress, winter
canola, Brussels sprout, carrot (GenBank Accession No.
CAB69453), Dutchman's breeches, spurge, daylily, winter
barley, Virginia waterleaf, narrow-leaved plantain,
30 plantain, speargrass, Kentucky bluegrass, Eastern
cottonwood, white oak, winter rye (Sidebottom et al., 2000,
Nature 406: 256), bittersweet nightshade, potato, chickweed,
dandelion, spring and winter wheat, triticale, periwinkle,
violet and grass.

The ISPs can be obtained by extraction from native sources by any suitable process, for example the isolation processes as described in WO 98/04699 and WO 98/4148.

5

Alternatively, ISPs can be obtained by the use of recombinant technology. For example host cells, typically micro-organisms or plant cells, may be modified to express ISPs and the ISPs may then be isolated and used in
10 accordance with the present invention. Techniques for introducing nucleic acid constructs encoding ISPs into host cells are well known in the art.

Typically, an appropriate host cell or organism would be
15 transformed by a nucleic acid construct that encodes the desired ISP. The nucleotide sequence coding for the polypeptide can be inserted into a suitable expression vector encoding the necessary elements for transcription and translation and in such a manner that they will be expressed
20 under appropriate conditions (e.g. in proper orientation and correct reading frame and with appropriate targeting and expression sequences). The methods required to construct these expression vectors are well known to those skilled in the art.

25

A number of expression systems may be used to express the polypeptide coding sequence. These include, but are not limited to, bacteria, fungi (including yeast), insect cell systems, plant cell culture systems and plants all
30 transformed with the appropriate expression vectors. Preferred hosts are those that are considered food grade - 'generally regarded as safe' (GRAS).

Suitable fungal species include yeasts such as (but not limited to) those of the genera *Saccharomyces*, *Kluyveromyces*, *Pichia*, *Hansenula*, *Candida*, *Schizosaccharomyces* and the like, and filamentous species such as (but not limited to) those of the genera *Aspergillus*, *Trichoderma*, *Mucor*, *Neurospora*, *Fusarium* and the like. Preferably the species selected is a yeast, most preferably a species of *Saccharomyces* such as *S. cerevisiae*. Where glycosylation of the ISP leads to reduced activity then it is preferred that the host exhibits reduced glycosylation of heterologous proteins.

A wide variety of plants and plant cell systems can also be transformed with the nucleic acid constructs of the desired polypeptides. Suitable plant species include maize, tomato, tobacco, carrots, strawberries, rape seed and sugar beet.

The sequences encoding the ISPs are preferably at least 80% identical at the amino acid level to an ISP identified in nature, more preferably at least 95% or 100% identical. However, persons skilled in the art may make conservative substitutions or other amino acid changes that do not reduce the RI activity of the ISP. For the purpose of the invention these ISPs possessing this high level of identity to an ISP that naturally occurs are also embraced within the term "ISPs".

Frozen confectionery products

Frozen confectionery products of the present invention comprise a plurality of discrete water ice confections. The water ice confections are not separated from one another by the use of wrappings or other non-edible packaging, or by compartmentalisation. Instead, the individual water ices

are packaged such that they are able to contact directly other individual frozen confections. However, the individual water ices are able to move relative to each other, in other words they are not immobilised within, for
5 example, a matrix such as a coating.

In a highly preferred embodiment, the water ice of the invention is free-flowing. Preferably, the frozen confectionery product of the invention remains free-flowing
10 after storage at -10°C for at least 10 days, more preferably at least 15 or 20 days.

The frozen confections are relatively small, for example having an average volume of less than 1 ml, more preferably
15 less than 0.5 ml. By way of example, beads having a diameter of from 5 mm to 10 mm would have a volume of from about 0.065 ml to about 0.5 ml. Typically, the discrete frozen confections have a minimum average volume such that each confection can be readily distinguished by a consumer.
20 For example, the discrete frozen confection preferably have a minimum average volume of at least about 0.02 ml.

The discrete frozen confections may be made to any shape, such as in the form of cubes or spheres. Preferably, the
25 frozen confections are substantially spherical.

The frozen confections may be in the form of a composite product where at least one portion or region of the product, such as a core or layer, does not contain ISPs. An example
30 of this would be a product containing a core of ice cream which lacks ISP, coated in a layer of water ice that does contain ISP. Preferably, substantially the outer layer of the composition confection comprises ISP, i.e. the region which will come into contact with other discrete frozen

confections. It will be appreciated that in the case of a composite product, the wt% amount of ISP added is calculated solely in relation to those components of the confection that contain ISP and not in relation to the complete
5 product.

Frozen confections may be aerated or unaerated. By unaerated is meant a frozen confection having an overrun of less than 20%, preferably less than 10%. An unaerated frozen
10 confection is not subjected to deliberate steps such as whipping to increase the gas content. Nonetheless, it will be appreciated that during the preparation of unaerated frozen confections, low levels of gas, such as air, may be incorporated in the product.

15 Water ice confections typically contain sugar, water, colour, fruit acid or other acidifying agent, fruit or fruit flavouring and stabiliser. Preferably, the total solids content is at least 6 wt%, more preferably at least 8 wt% or
20 at least 10, 12, 15 or 20 wt% and may be as high as about 35 wt%. Preferably the total solids content is less than 35 wt%, more preferably less than 25 wt%. Water ices may be aerated or unaerated. If aerated, the overrun is typically less than about 50%, for example from about 25% to 30%. In
25 one embodiment, the water ice confections of the invention are unaerated.

Preferably the water ice confections contain less than 2 wt% artificial sweeteners, more preferably less than 1 wt%. In
30 a highly preferred embodiment, no artificial sweeteners, such as aspartame or acesulfame are present in the water ice confections.

Frozen confections of the invention typically comprise one or more stabiliser, such as one or more stabilisers selected from gums, agar, alginates and derivatives thereof, gelatin, pectin, lecithin, sodium carboxymethylcellulose, carrageenan and furcelleran. Preferably a blend of stabilisers is used, such as blend of a gum and carrageenan. In a preferred embodiment, the frozen confection comprises from 0.1 to 1 wt% stabiliser.

10 Frozen confections of the invention typically comprise at least about 0.0005 wt% ISP. ISPs can be used at very low concentrations and therefore preferably the confections comprise less than 0.05 wt% ISP. A preferred range is from about 0.001 to 0.01 wt%.

15 Frozen confections of the invention can be manufactured using a number of techniques known in the art. For example, free-flowing beads can be manufactured by dispensing drops of the liquid mix into a freezing chamber of liquid nitrogen (see WO96/29896). Other shapes can be manufactured by moulding techniques, for example by introducing a liquid premix into a cooled mould. Moulded products may contain complex shapes and have a high degree of surface definition.

25 Ice cream-containing products and the like need not be subjected to a cold hardening step of below from -20°C to -25°C, although this may be used if desired, especially if the product is a composite product with a layer or core that does not contain ISP.

30 The frozen confectionery product of the invention may be packaged in containers for sale to consumers as an individual unit. The volume of such containers is typically from 100 ml to 1000 ml, such as from 200 ml to 500 ml.

However, the product can also be packaged in larger containers for retail purposes where the product is dispensed into smaller containers at the retail premises, e.g. in fast food outlets or as a pick 'n' mix format where consumers can choose from frozen confections of the invention having different shapes, flavours and/or colours. These larger containers may, for example, have a volume greater than about 1000 ml, for example at least 2000 ml or 5000 ml.

10

The present invention will now be further described with reference to the following examples, which are illustrative only and non-limiting.

15 **EXAMPLES**

Examples 1 to 5 and Comparative Examples 1 to 4
- water ice beads

20 **Materials and methods**

Water ice premixes were produced according to the following recipes.

Table 1

25

Ingredients	C. Ex. 1	Ex. 1a	Ex. 1b	Ex. 1c	Ex. 1d
Sugar source (I)	15.0	15.0	15.0	15.0	15.0
Stabiliser (II)	0.35	0.35	0.35	0.35	0.35
Colour (III)	0.088	0.088	0.088	0.088	0.088
Flavouring (IV)	0.31	0.31	0.31	0.31	0.31
Fat source (V)	0.8	0.8	0.8	0.8	0.8
Emulsifier (VI)	0.2	0.2	0.2	0.2	0.2
Fruit juice concentrate (VII)	5.0	5.0	5.0	5.0	5.0
Food acid (VIII)	0.32	0.32	0.32	0.32	0.32
Water	77.929	78	78.4	78.98	79.4

13

ISP (%)	0	0.0005	0.0025	0.005	0.007
Fat (%)	1.0	1.0	1.0	1.0	1.0
Total solids (%)	20	20	20	20	20

Ingredients	C. Ex. 2	Ex. 2	C. Ex. 3	Ex. 3
Sugar source (I)	13.7	13.7	14.0	14.0
Stabiliser (II)	0.353	0.353	0.353	0.353
Artificial sweetener (VIV)	0	0	0	0
Colour (III)	0.088	0.088	0.088	0.088
Flavouring (IV)	0.31	0.31	0.31	0.31
Fat source (V)	0.8	0.8	0.8	0.8
Emulsifier (VI)	0.2	0.2	0.2	0.2
Salt	0	0	0	0
Fruit juice concentrate (VII)	5.0	5.0	5.0	5.0
Food acid (VIII)	0.32	0.32	0.32	0.32
Water	79.229	80.8	78.929	80.8

ISP (%)	0	0.005	0	0.005
Fat (%)	1.0	1.0	1.0	1.0
Total solids (%)	15	15	9	9

Ingredients	C. Ex. 4	Ex. 4	Ex. 5
Sugar source (I)	4.21	4.21	15.5
Stabiliser (II)	0.35	0.35	0.35
Artificial sweetener (VIV)	0.036	0.036	0
Colour (III)	0.088	0.088	0.11
Flavouring (IV)	0.31	0.31	0.40
Fat source (II)	0.8	0.8	0.8
Emulsifier (VI)	0.2	0.2	0.2
Salt	0.09	0.09	0
Fruit juice concentrate (VII)	0	0	5.2
Food acid (VIII)	0.32	0.32	0.77
Water	93.59	93.59	75.6

ISP (%)	0	0.005	0.005
Fat (%)	1.0	1.0	1.0
Total solids (%)	6	6	20

5

I Sugar source can be any typically used water ice ingredient such as either sucrose or fructose or a blend of sucrose/fructose in 97/3 ratio or sucrose/fructose in 54/46 ratio.

10 II A blend of pectin/carrageenan.

III Any typically used water ice colour.

- IV Any typically used water ice flavourings.
- V Fat source such as coconut oil or other bland fat type.
- VI Emulsifier such as monoglycerolpalmitate (MGP).
- VII Fruit juice concentrate added to give flavour/fruit
5 value, solids should be balanced if added: level shown
is an example and can be any fruit
- VIII Any typically used water ice food acid such as citric
acid.
- VIV Any typically used water ice artificial sweetener such
10 as acesulfame or aspartame or a 50/50 blend of both.

TS indicates the total solids content as a percentage by
weight.

TF indicates the total fat content (including emulsifier) as
a percentage by weight.

- 15 The determination of these values is conventional in the art

Mix process

- All dry ingredients with the exception of the acidifier mix
20 were added to water which was pre-heated to 80°C, followed
by stirring for 5 minutes. Then all the liquid ingredients
and acidifier were added, stored for 1 minute, pasteurised
at 82°C for 33 seconds, homogenised at 150-170 bar pressure
and cooled to 5°C until required. Glacein was added post
25 pasteurisation for the purposes of this study, addition pre-
pasteurisation would require removal of an equal weight of
water from the formulation.

Particle formation

- 30 The liquid mix at 5°C was loaded into a mix chamber of 5
litres capacity which fed directly into a dripping nozzle of

15

1 mm internal diameter. The liquid drops in turn fell into liquid nitrogen where they were rapidly frozen into approximately spherical balls. From here they were filled into a cylindrical type cup (height 95 cm, bottom outside diameter 63 cm, top outside diameter 46mm) to a fill weight of 85 g, from the base, the base being sealed on with an iron. The products were then placed at -25°C until required for measurement.

10 *Free flow test*

Samples are held at a constant temperature of either -10°C or -25°C for 50 days. Samples in a pot (six replicates) were squeezed manually at -25°C, the pot was then opened and upturned and the flow properties of the contents assessed on a 5 point scale according to which:

- 1 = particles exit pot and are completely free flowing.
- 2 = if particles do not exit at 1, pot is re-closed and inverted 5 times to separate the particles, which exit when the lid is opened and upturned.
- 3 = as 2 but two gentle squeezes to the sides are additionally required before particles will exit. No residual deformation of the pack is seen.
- 4 = as 3 but two harder squeezes are required which will deform the pack, leaving it still deformed after the particles are removed.
- 5 = particles can not be made to exit.

A squeeze score of 3 is considered the maximum in terms of acceptable flowability. The scores quoted in Table 2 are mean values of the scores obtained for six replicate samples. The test was performed with respect to time, sampling every few days.

Results

Table 2

5

Time (Days)	C. Ex. 1 Squeeze value		Ex. 1a Squeeze value		Ex. 1b Squeeze value		Ex. 1c Squeeze value		Ex. 1d Squeeze value	
	-10°C	-25°C	-10°C	-25°C	-10°C	-25°C	-10°C	-25°C	-10°C	-25°C
1	2	n.d	2	n.d	n.d	n.d	3	2	n.d	n.d
2	n.d	n.d	n.d	n.d	3	2	n.d	n.d	n.d	n.d
3	4	2	3	2	3	2	n.d	n.d	3	2
4	n.d	2	3	2	3	2	3	2	3	2
5	n.d	n.d	n.d	n.d	3	2	3	2	3	2
7	3	2	3	2	3	2	n.d	n.d	3	2
10	4	2	4	3	3	2	3	3	3	2
15	4	2	4	3	3	2	3	3	3	2
21	5	3	3	3	3	2	3	3	3	2
30	5	3	3	3	3	2	4	2	3	2
40	5	3	3	3	4	2	3	2	3	2
50	5	3	5	3	4	2	4	3	3	2

Time (Days)	C. Ex. 2 Squeeze value		Ex. 2 Squeeze value		C. Ex. 3 Squeeze value		Ex. 3 Squeeze value	
	-10°C	-25°C	-10°C	-25°C	-10°C	-25°C	-10°C	-25°C
1	3	1	2	1	3	2	2	1
2	n.d	n.d	2	1	3	2	2	1
3	n.d	n.d	2	2	3	2	2	2
4	5	2	n.d	n.d	n.d	n.d	n.d	n.d
5	4	2	n.d	n.d	n.d	n.d	n.d	n.d
7	4	2	3	2	3	3	3	2
10	4	2	3	3	4	2	3	2
15	4	2	3	2	4	3	3	2
21	5	2	3	2	4	3	3	2
30	5	3	3	2	4	3	3	2
40	5	3	3	2	5	3	3	2
50	5	3	3	2	5	3	3	3

Time (Days)	C. Ex. 4 Squeeze value		Ex. 4 Squeeze value		Ex. 5 Squeeze value	
	-10°C	-25°C	-10°C	-25°C	-10°C	-25°C
1	3	1	3	1	3	2
2	3	2	2	1	n.d	n.d
3	n.d	n.d	3	1	n.d	n.d
4	n.d	n.d	n.d	n.d	2	2
5	3	2	n.d	n.d	n.d	n.d
7	3	2	3	1	n.d	n.d

17

10	3	1	2	1	n.d	n.d
15	3	2	3	2	3	2
21	3	2	3	2	3	2
30	3	3	3	2	3	2
40	4	3	3	3	3	2
50	4	3	3	3	3	2
90	n.d	n.d	n.d	n.d	4	2

Comparative Example 1 is a control sample at 20% TS, which does not contain ISP. After 50 days at -25°C, the sample remained free flowing. After 10 days at -10°C the sample became unacceptable.

Example 1a contains 0.0005% ISP. After 50 days -25°C, the sample remained free flowing. After 50 days at -10°C, the sample became unacceptable.

Example 1b contains 0.0025% ISP. After 50 days at -25°C, the sample remained free flowing. After 40 days at -10°C, the sample became unacceptable.

Example 1c contains 0.005% ISP. After 50 days at -25°C, the sample remained free flowing. After 50 days at -10°C, the sample became unacceptable.

Example 1d contains 0.007% ISP. The sample remained free flowing throughout the test at both -25°C and -10°C. This sample showed marked improvement over comparative example 1 and examples 1a, 1b, and 1c.

Comparative Example 2 is a control sample at 15% TS, which does not contain ISP. After 50 days at -25°C, the sample remained free flowing. After 4 days at -10°C, the sample became unacceptable.

Example 2 contains 0.005% ISP at 15% TS. The sample remained free flowing throughout the test at -25°C and -10°C.

5

Comparative Example 3 is a control sample at 9% TS, which does not contain ISP. After 50 days at -25°C, the sample remained free flowing. After 10 days at -10°C, the sample became unacceptable.

10

Example 3 contains 0.005% ISP at 9% TS. The sample remained free flowing throughout the test at both -25°C and -10°C.

Comparative Example 4 is a control sample at 6% TS, which does not contain ISP. After 50 days, the sample remained free flowing. After 40 days at -10°C, the sample became unacceptable.

Example 4 contains 0.005% ISP at 6% TS. The sample remained free flowing throughout the test at both -25°C and -10°C.

Example 5 contains 0.005% ISP at 20% TS. After 90 days at -25°C, the sample remained free flowing. After 90 days at -10°C, the sample became unacceptable, showing marked improvement over comparative example 1.

All publications mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described methods and products of the invention will be apparent to those skilled in the art without departing from the scope of the invention. Although the invention has been described in connection with specific

preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are
s apparent to those skilled in the relevant fields are intended to be within the scope of the following claims.